

AD-A205 487

DTIC FILE COPY

(4)

OFFICE OF NAVAL RESEARCH

Contract N00014-86-K-0043

TECHNICAL REPORT No. 91

Decay of Molecules at Corrugated Thin Metal Films

by

P. T. Leung, Young Sik Kim and Thomas F. George

Prepared for Publication

in

Physical Review B

Departments of Chemistry and Physics  
State University of New York at Buffalo  
Buffalo, New York 14260

February 1989

Reproduction in whole or in part is permitted for any purpose of the  
United States Government.

This document has been approved for public release and sale;  
its distribution is unlimited.

DTIC  
ELECTE  
S 8 FEB 1989 D  
CE

89

2

28

010

UNCLASSIFIED  
SECURITY CLASSIFICATION OF THIS PAGE


REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution unlimited		
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) UBUFFALO/DC/89/TR-91			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Depts. Chemistry & Physics State University of New York		6b. OFFICE SYMBOL (if applicable)	7a. NAME OF MONITORING ORGANIZATION		
6c. ADDRESS (City, State, and ZIP Code) Fronczak Hall, Amherst Campus Buffalo, New York 14260		7b. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217			
8a. NAME OF FUNDING / SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (if applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER Contract N00014-86-K-0043		
8c. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217		10. SOURCE OF FUNDING NUMBERS			
		PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.	WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) Decay of Molecules at Corrugated Thin Metal Films					
12. PERSONAL AUTHOR(S) P. I. Leung, Young Sik Kim and Thomas F. George					
13a. TYPE OF REPORT		13b. TIME COVERED FROM TO		14. DATE OF REPORT (Year, Month, Day) February 1989	
				15. PAGE COUNT 27	
16. SUPPLEMENTARY NOTATION Prepared for publication in Physical Review B					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP			
			MOLECULAR DECAY / SURFACE PLASMONS /		
			THIN METAL FILMS / LONG- AND SHORT-RANGE		
			CORRUGATED CROSS-COUPLING		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) The problem of the decay rate for molecules at corrugated thin metal films is considered theoretically via a classical phenomenological approach by generalizing a previous theory for a single rough surface. In particular, numerical results are worked out in detail for grating films with various possible arrangements of the two grating surfaces, and for both the cases with a supported and free-standing film. Effects due to cross-coupling into the long- and short-range surface plasmons are observed, and correlations to recent experiments performed by the Rochester group are made. It is suggested that fluorescence experiments with pulsed-laser excitations should be carried out in order for the results in the present calculation to be observed.					
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED / UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. David L. Nelson			22b. TELEPHONE (Include Area Code) (202) 696-4410		22c. OFFICE SYMBOL

DD Form 1473, JUN 86

Previous editions are obsolete.

SECURITY CLASSIFICATION OF THIS PAGE

UNCLASSIFIED



Decay of Molecules at Corrugated Thin Metal Films

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

P. T. Leung  
Department of Physics  
Portland State University  
P. O. Box 751  
Portland, Oregon 97201

Young Sik Kim\* and Thomas F. George  
Departments of Chemistry and Physics & Astronomy  
239 Fronczak Hall  
State University of New York at Buffalo  
Amherst, New York 14260

Abstract

The problem of the decay rate for molecules at corrugated thin metal films is considered theoretically via a classical phenomenological approach by generalizing a previous theory for a single rough surface. In particular, numerical results are worked out in detail for grating films with various possible arrangements of the two grating surfaces, and for both the cases with a supported and a free-standing film. Effects due to cross-coupling into the long- and short-range surface plasmons are observed, and correlations to recent experiments performed by the Rochester group are made. It is suggested that fluorescence experiments with pulsed-laser excitations should be carried out in order for the results in the present calculation to be observed.

\* Present address: Department of Chemistry, Princeton University,  
Princeton, New Jersey 08544

PACS Nos: 73.60.Ag, 73.90.+f, 78.65.-s, 82.65.Pa

## I. Introduction

The problem of determining lifetimes for molecules in the proximity of metal surfaces has been investigated for the past twenty years due to its fundamental and practical importance.<sup>1</sup> Both the cases of a flat<sup>2</sup> and a structured<sup>3-5</sup> surface have been thoroughly studied, though it seems to the present authors that for the latter type of surfaces, more experimental studies are needed to check the various results obtained from theoretical modeling.<sup>2-5</sup> Related to, but in itself a more interesting problem, is the case where the ad molecules are placed close to a thin-film substrate. Roughly speaking, when the thickness is thin enough, a thin-film system is no more than two surfaces interferring with each other. For the case of a flat film, both the problems of molecular fluorescence lifetimes<sup>2,6</sup> and light-scattering by an external source<sup>7</sup> have been studied, with the latter having been extended to optically nonlinear films.<sup>8</sup> A system of perhaps even greater interest is a corrugated thin film, since the two surface plasmons supported by the two sides can now couple with each other leading to much richer proximity effects. This cross-coupling effect has been studied recently by means of light-scattering experiments in which effects such as the couplings into long- and short-range surface plasmons<sup>9-16</sup> and the dramatic enhanced transparency<sup>17,18</sup> of a metallic film have been observed. The problem of molecular fluorescence in the vicinity of such a system has, however, received less attention. Only until very recently, being stimulated by a controversy among different groups as to whether cross-coupling has really been observed from the asymmetry of the reflectance dips,<sup>12</sup> a group at the University of Rochester has carried out successive experiments<sup>19-21</sup> to study the fluorescence of photoresist molecules in the vicinity of a Ag grating film, and it is claimed that cross-coupling has been unambiguously observed in their experiments.

Motivated by the Rochester experiments, we have set forth here to perform a theoretical investigation of the decay rates (inverse lifetimes) for molecules near a metallic grating film by extending our previous theory for a grating surface.<sup>5</sup> We shall consider different structures of the system, including both the asymmetrically and symmetrically (free-standing) bound cases and films with only one surface roughened, and we shall make correlations with previous light-scattering<sup>9-16</sup> or fluorescence studies<sup>19-21</sup> wherever possible. In the following, we shall present in Sec. II a phenomenological theory, and numerical illustrations of various decay-rate spectra will follow in Sec. III. In Sec. IV we shall discuss the possible experimental implications from our calculations.

Before we present our formalism, we want to emphasize that whereas the theoretical treatment of the light-scattering problem usually incorporates all the dynamics of the problem,<sup>10,11</sup> the case for the calculation of the decay rate problem in the vicinity of a rough surface has been simplified to work in the nonretarded (static) limit.<sup>3,4</sup> We have reevaluated the applicability of the nonretarded approach in terms of the substrate conductivity and the molecule-surface distances, and have emphasized the significance of applying a dynamic theory for such decay rate calculations.<sup>1</sup> Very recently, we have shown further that the morphology-induced resonances in the decay rate spectrum can only occur in the dynamic treatment but not in the static approach, due to the neglect of radiative energy transfer in the static theory.<sup>22,23</sup> Hence, we propose that to treat the cross-coupling effects (mediated by the grating period) in the decay-rate spectrum for the admolecules, it is necessary to impose a dynamic theory for the description, which we now proceed to develop in the following section.

## II. FORMALISM

The geometry of our problem is depicted in Fig. 1, where we consider a two-level system (modeled by a dipole  $\vec{\mu}$ ) located at a distance  $z = d$  above a thin metal (taken as Ag) film bounded by two grating surfaces located at  $z = 0$  and  $z = -t$ , respectively. Thus the profile functions take the simple forms

$$\zeta_1 = \zeta_{o1} e^{iQ_1 x}, \quad \zeta_2 = \zeta_{o2} e^{iQ_2 x}, \quad (1)$$

respectively, where we shall assume small corrugations ( $\zeta_{oi} Q_i \ll 1$ ) so that perturbation theory can be applied.<sup>5</sup> The optical properties of the three media are described by their dielectric functions,  $\epsilon_1$ ,  $\epsilon_2 = \epsilon'(\omega) + i\epsilon''(\omega)$  and  $\epsilon_3$ . According to the classical phenomenological (CP) approach, we must calculate the reflected field ( $\vec{E}^r$ ) acting on the emitting dipole, which is then the cause of the induced decay, and we then obtain the normalized decay rate as<sup>2</sup>

$$\frac{\gamma}{\gamma_0} = 1 + \frac{3q\epsilon_1}{2k_1^3} \text{Im}G(\omega), \quad (2)$$

with

$$G(\omega) = \frac{\vec{E}^r(\omega)}{\mu}, \quad (3)$$

and  $q$  being the quantum yield of the emitting state,  $k_1 = \frac{\sqrt{\epsilon_1}\omega}{c}$  the emission wave number and  $\gamma_0$  the decay rate for a free molecule. Since the details of similar calculations for a single grating surface have been given in a

previous article,<sup>5</sup> here we shall only sketch briefly the important steps in obtaining our final results. where as before, we shall consider the simple case with the dipole oriented perpendicular to the film.

By generalizing the original integral-equation theory for light-scattering at a corrugated surface,<sup>24</sup> Mills and Maradudin have obtained for the corrugated film the  $\mu$ -th component for the roughness contribution to  $\vec{E}^r$  to first-order perturbation, given by<sup>25</sup>

$$\begin{aligned}
 E_{\mu}^r(\vec{r};\omega) = & - \frac{k^2}{16\pi^3} \int d^2k_{\parallel} e^{i\vec{k}_{\parallel} \cdot \vec{r}_{\parallel}} \int dz' d_{\mu\nu}(\vec{k}_{\parallel}|\omega|zz') ([\epsilon_2\theta(z'+t) - \epsilon_1] \\
 & \times \delta(z')\hat{f}_1(\vec{k}_{\parallel}-\vec{k}_{\parallel}^{(0)}) + [\epsilon_3-\epsilon_2\theta(-z')] \delta(z'+t) \\
 & \times [\hat{f}_2(\vec{k}_{\parallel}-\vec{k}_{\parallel}^{(0)}) + t]) E_{\nu}^{(0)}(\vec{k}_{\parallel}^{(0)}|\omega|z') , \quad (4)
 \end{aligned}$$

where  $\parallel$  denotes any vector on the xy-plane,  $\hat{f}_1$  and  $\hat{f}_2$  are Fourier transforms of Eq. (1),  $\theta$  is the Heaviside step function, and  $d_{\mu\nu}$  is the Fourier transform of the two-dimensional "flat propagator" obtained in Ref. 25.  $E_{\nu}^{(0)}$  in Eq. (4) denotes the total field for the dipole-substrate problem in the medium that contains  $z'$  for the case of a perfectly flat film, and expressions which, in terms of the Green dyadic,<sup>26</sup> have already been worked out in the literature by Chance, Prock and Silbey.<sup>2</sup> As pointed out before,<sup>5</sup> to evaluate the integral of  $\delta(z')$  in Eq. (4) where  $E_{\nu}^{(0)}$  may be discontinuous across the boundary of the films, one must adopt Agarwal's modifications<sup>27</sup> and not just take the mean value of the integrals at each side of the boundary.<sup>24</sup> It is straightforward to show that Agarwal's modifications to the theory of Maradudin and Mills<sup>24</sup> should be applicable to multilayered systems as well. Adopting further the

normal incidence approximation ( $\vec{k}_{\parallel}^{(0)} \approx 0$ ), which is good for shallow gratings,<sup>5</sup> we finally obtain the roughness contribution to the reflected field at the dipole site, given by ( $k = \frac{\omega}{c}$ )

$$E_z^r(d; \omega) = - \frac{k^2}{4\pi} [(\epsilon_2 - \epsilon_1) \zeta_{o_1} I_1 + (\epsilon_3 - \epsilon_2) \zeta_{o_2} I_2] , \quad (5)$$

where  $I_1$  and  $I_2$  are integrals given by:

$$I_1 = \frac{4\pi Q_1^2}{\alpha_1 k^2 \epsilon_1^2} e^{i\alpha_1 d} \frac{\left(\frac{\epsilon_2}{\epsilon_3} + \frac{\alpha_2}{\alpha_3}\right) e^{i\alpha_2 t} - \left(\frac{\epsilon_2}{\epsilon_3} - \frac{\alpha_2}{\alpha_3}\right) e^{-i\alpha_2 t}}{\left(\frac{\alpha_2}{\alpha_1} - \frac{\epsilon_2}{\epsilon_1}\right) \left(\frac{\alpha_2}{\alpha_3} + \frac{\epsilon_2}{\epsilon_1}\right) e^{i\alpha_2 t} - \left(\frac{\alpha_2}{\alpha_1} + \frac{\epsilon_2}{\epsilon_1}\right) \left(\frac{\alpha_2}{\alpha_3} - \frac{\epsilon_2}{\epsilon_1}\right) e^{-i\alpha_2 t}} \times \int_0^\infty d\lambda \frac{\lambda^3}{h_1} e^{ih_1 d} f_1 , \quad (6)$$

$$I_2 = \frac{4\pi Q_2^2}{\beta_1 k^2 \epsilon_1^2} \left(\frac{k_1}{k_3}\right) e^{i\beta_1 d} \frac{2\beta_2/\beta_3}{\left(\frac{\beta_2}{\beta_1} - \frac{\epsilon_2}{\epsilon_1}\right) \left(\frac{\beta_2}{\beta_3} + \frac{\epsilon_2}{\epsilon_1}\right) e^{i\beta_2 t} - \left(\frac{\beta_2}{\beta_1} + \frac{\epsilon_2}{\epsilon_1}\right) \left(\frac{\beta_2}{\beta_3} - \frac{\epsilon_2}{\epsilon_1}\right) e^{-i\beta_2 t}} \times \int_0^\infty d\lambda \frac{\lambda^3}{h_1} e^{ih_2 d} f_2 , \quad (7)$$

where

$$\alpha_i^2 = \epsilon_i \frac{\omega^2}{c^2} - Q_1^2 , \quad \beta_i^2 = \epsilon_i \frac{\omega^2}{c^2} - Q_2^2 , \quad (8)$$

and



$$h_i^2 = k_i^2 - \lambda^2 \quad \text{with} \quad k_i = \sqrt{\epsilon_i} \frac{\omega}{c} \quad (9)$$

Note that the signs of  $\alpha_i$  and  $\beta_i$  must be carefully assigned so as to satisfy the boundary conditions for the fields at infinity.<sup>26</sup> The functions  $f_1$  and  $f_2$  are defined as<sup>2</sup>

$$f_1 = \frac{2Sh_1\epsilon_2}{Sh_1\epsilon_2 + h_2\epsilon_1} \quad (10)$$

with

$$S = \frac{h_2\epsilon_3 - ih_3\epsilon_2 \tan(h_2\tau)}{h_3\epsilon_2 - ih_2\epsilon_3 \tan(h_2\tau)} \quad (11)$$

and

$$f_2 = \frac{k_2}{k_3} (f_3 e^{ih_2\tau} + f_4 e^{-ih_2\tau}) \quad (12)$$

with

$$f_3 = \frac{1}{2} \left( \frac{h_1 k_2}{h_2 k_1} + \frac{k_1}{k_2} \right) - \frac{f_1 - 1}{2} \left( \frac{h_1 k_2}{h_2 k_1} - \frac{k_1}{k_2} \right) \quad (13)$$

$$f_4 = \frac{1}{2} \left( \frac{k_1}{k_2} - \frac{h_1 k_2}{h_2 k_1} \right) + \frac{f_1 - 1}{2} \left( \frac{k_1}{k_2} + \frac{h_1 k_2}{h_2 k_1} \right) \quad (14)$$

Hence, from Eqs. (2) and (3), we finally obtain to first order in roughness the reduced total decay rate for the admolecule, given by (assuming  $q = 1$ )

$$\frac{\gamma}{\gamma_0} = \gamma^F + \gamma^R, \quad (15)$$

with

$$\gamma^F = 1 + \frac{3\epsilon_1}{2k_1^3} \text{Im}G^F, \quad (16)$$

$$\gamma^R = \frac{3\epsilon_1}{2k_1^3} \text{Im}G^R, \quad (17)$$

where

$$G^F(\omega) = \frac{i}{\epsilon_1} \int_0^\infty d\lambda \frac{\lambda^3}{h_1} (f_1 - 1) e^{2ih_1 d}, \quad (18)$$

$$G^R(\omega) = \frac{E_z^R(d; \omega)}{\mu}, \quad (19)$$

and  $E_z^R(d; \omega)$  as given in Eq. (5). The correctness of this result can be checked by taking various limits such as by letting one of the  $\epsilon_{oi}$ 's go to zero and by making any of the two neighboring  $\epsilon$ 's to be identical, and the limiting results can then be checked with those obtained previously for a single grating surface.<sup>5</sup>

### III. NUMERICAL RESULTS

Here we apply our general results in Sec. II to several film systems of interest, each of which has been studied previously in the literature for light-scattering properties.

#### A. Film with Only One Grating Surface

This system has recently been studied by Friar and coworkers<sup>15</sup> for the reflectance from an impinging plane wave. To investigate our decay-rate problem, we simply set  $\zeta_{o1} = 0$  in Eq. (1) and are hence left with only the term  $\zeta_{o2} I_2$  in Eq. (5). Furthermore, we take  $\epsilon_1 = 1.0$  and  $\epsilon_3 = 3.6$ , and  $\epsilon_2 = \epsilon' + i\epsilon''$  for silver can be found from various source books.<sup>28</sup> Other parameters are as given in the figure captions.

Figure 2 shows the total decay ( $\gamma$ , solid line) as well as the separate contributions from the flat ( $\gamma^F$ , dotted line) and rough ( $\gamma^R$ , dashed line) components of the film. Here various resonance peaks can be clearly observed corresponding to the single-surface nonradiative transfer ( $\sim 3.6$  eV),<sup>5</sup> the flat-film interference resonance,<sup>2</sup> and the morphology induced resonance<sup>22</sup> due to the grating surface, respectively.

Figure 3 shows, for the same system, the total rate  $\gamma$  for different thicknesses of the film. It is obvious that as  $t \rightarrow \infty$ , the single flat surface result (dotted line) is recovered, where both the flat-film interference and grating-induced resonance disappear. Similar effects on the light-scattering resonances due to the variation of film thickness in this system have been observed in Ref. 15. Likewise, similar results are obtained if we set  $\zeta_{o2} = 0$  but  $\zeta_{o1} \neq 0$  in our expressions in Sec. II,<sup>14</sup> except that, of course, in the limit  $t \rightarrow \infty$ , the grating induced peak ( $\sim 2$  eV) still shows up, in agreement with our previous investigations.<sup>22</sup>

### B. Supported Grating Film

Figure 4 shows results for  $\gamma$ ,  $\gamma^F$  and  $\gamma^R$ , where we have assumed  $\zeta_{o1} = \zeta_{o2} = 5 \text{ \AA}$ . This asymmetrically-bound system ( $\epsilon_1 = 1$ ,  $\epsilon_3 = 3.6$ ) is known for its "inefficiency" of cross-coupling between the two surface plasmons supported by each grating surface.<sup>21</sup> In fact, the resonance peaks that appear in Figs. 4 and 5 are not much different (except in strength) when compared with the results in case A. This fact can be easily understood in the case of plane-wave incidence, where the dispersion relations for the two surface plasmons are simpler, from which serious restrictions are found to be imposed on the coupling condition when the film is bounded by two different dielectric media.

### C. Free-Standing Grating Film

This is the most efficient system for cross-coupling studies, since the two surface plasmons are coupled for all energies in the case of plane-wave incidence.<sup>21</sup> The fabrication of such a film has been described by Inagaki and coworkers.<sup>13</sup> For simplicity, we have assumed  $\epsilon_1 = \epsilon_3 = 1$  and  $\zeta_{o1} = \zeta_{o2} = 5 \text{ \AA}$ . It is clearly seen from Fig. 6 that the two surface plasmons on each grating surface are coupled to the long- ( $\omega \sim 1.5 \text{ eV}$ ) and short-range ( $\omega \sim 2.0 \text{ eV}$ ) surface plasmons. Similar features have been observed in a light-scattering experiment<sup>13</sup> in which the photoacoustic technique has been applied to monitor the excitation of such cross-coupled plasmons. However, the splitting of higher-order peaks observed there does not show up in our calculation due to the different dispersion relations which we have here for the coupling of the dipole radiations to the film, instead of the plane-wave couplings as in the light-scattering case, where one has multiple branches of dispersion relations.<sup>13</sup> Figure 7 shows results for the same system with different film thicknesses, where again the cross-coupling effect disappears for a thick film, just as in the experiments of Inagaki and coworkers.<sup>13</sup>

#### D. Remarks on the Rochester Experiments

In three recent investigations,<sup>19-21</sup> a group at the University of Rochester has performed beautiful experiments in which cross-coupling of the two surface plasmons from the sides of a thin grating Ag film is unambiguously observed, through the scattered (transmitted) light from the molecular fluorescence of a photoresist in the vicinity of the film. Both ~~asymmetrically~~- and symmetrically-bound films have been studied. We have performed a calculation of the decay rates [Eqs. (5)-(19)] for one of their systems using their parameters for the optical constants and the silver grating of different periods, and our results are shown in Fig. 8. Although we observe a similar qualitative feature to that in Fig. 1 of Ref. 19, the positions of the peaks are quite different in the two cases. We suspect that the type of experiments we are referring to would be quite different from the Rochester experiments, since in the latter the dispersion relations are still guided by those derived from an incident plane wave, and the fluorescent molecules are simply undergoing Thomson scattering by which the incident light from the optical fiber is transmitted through the grating film. Here, as discussed above, we are instead referring to the coupling between the surface plasmons and the dipole fields of the decaying molecules in which a superposition of all the harmonics are involved, and hence we expect the resonances to be governed by different dispersion relations. As a matter of fact, one of the Rochester experiments<sup>21</sup> indeed shows that for the symmetrically-bound case, multiple resonances due to higher-order dispersions do appear in their transmittance spectrum, much as in the case of plane-wave scattering as studied by Inagaki et al.<sup>13</sup> In fact, the peaks shown here for the decay-rate spectrum in Fig. 8 are not much different from those due to single-surface (thick-film) couplings (cf. Figs. 3 and 5). Cross-coupling

into the split modes of the long- and short-range surface plasmons can only be seen here if the film thickness is further decreased, as shown in Fig. 9.

We would like to propose that for the observation of the peaks calculated in this present work, fluorescence experiments have to be performed in which the molecules are excited by a short-pulse laser and the emission intensity is monitored by various time-resolved techniques.<sup>29</sup> In this fashion, the decay rate values can then be deduced by assuming appropriate forms for the decay curves that fit the intensity data. In such experiments, the pulse width of the laser must be considerably shorter than the lifetimes ( $\sim 1/\gamma$ ) of the admolecules, and the molecule decaying from its excited state (after being "pumped up" by the laser pulse) can then be modeled as a radiating dipole.<sup>2</sup> The reflected field from the substrate film acting on the dipole (which leads to the induced decay of the molecule) can then be correctly described by the present formalism, and hence the peak positions from the present work can be checked by exciting various fluorescent molecules of different natural frequencies.

#### IV. CONCLUSION

We have performed calculations of the decay-rate spectrum for fluorescent molecules near a corrugated metal film for both supported and unsupported systems, and appropriate types of experiments have been suggested. While many qualitative features observed here are similar to the case of light-scattering analyses of such film systems,<sup>9-16</sup> we emphasize that a distinct feature of the present problem is the dipole nature of the radiation coupled to the surface plasmons of the film. Because of the difference between this coupling and the plane-wave coupling to the grating modes, we expect that interesting phenomena will arise if one studies the

photoabsorption of these admolecules in which an external laser field (plane wave) is introduced to drive the admolecules with both the external and molecular (dipolar) fields coupled to the surface plasmon modes at the same time. We have obtained some preliminary results for such a problem, and details will be presented in a separate paper.<sup>30</sup>

#### ACKNOWLEDGMENTS

This research was supported by the Office of Naval Research, the National Science Foundation under Grant CHE-8620274 and the Air Force Office of Scientific Research (AFSC), United States Air Force, under Contract F49620-86-C-0009. The United States Government is authorized to reproduce and distribute reprints for governmental purposes notwithstanding any copyright notation hereon.

# REFERENCES

1. For a review, see P. T. Leung and T. F. George, *Spectroscopy* 4, 35 (1989); H. Metiu, *Prog. Surf. Sci.* 17, 153 (1984).
2. See the comprehensive review by R. R. Chance, A. Prock and R. Silbey, *Adv. Chem. Phys.* 37, 1 (1978).
3. J. Arias, P. K. Aravind and H. Metiu, *Chem. Phys. Lett.* 85, 404 (1982).
4. P. T. Leung, Z. C. Wu, D. A. Jelski and T. F. George, *Phys. Rev. B* 36, 1475 (1987).
5. P. T. Leung and T. F. George, *Phys. Rev. B* 36, 4664 (1987).
6. See also W. R. Holland and D. G. Hall, *Opt. Lett.* 10, 414 (1985).
7. See, e.g., E. N. Economou, *Phys. Rev.* 182, 539 (1969); A. B. Buckman and C. Kuo, *J. Opt. Soc. Am.* 69, 343 (1979).
8. See, e.g., W. Chen and D. L. Mills, *Phys. Rev. B* 35, 524 (1987); W. Chen and A. A. Maradudin, *J. Opt. Soc. Am. B* 5, 529 (1988).
9. I. Pockrand, *Opt. Commun.* 13, 311 (1975); I. Pockrand and H. Raether, *Appl. Opt.* 16, 1784 (1977).
10. D. Sarid, *Phys. Rev. Lett.* 47, 1927 (1981).
11. G. S. Agarwal, *Phys. Rev. B* 31, 3534 (1985); S. D. Gupta, G. V. Varada and G. S. Agarwal, *Phys. Rev. B* 36, 6331 (1987).
12. M. G. Weber and D. L. Mills, *Phys. Rev. B* 32, 5057 (1985).
13. T. Inagaki, M. Motosuga, E. T. Arakawa and J. P. Goudonnet, *Phys. Rev. B* 31, 2548 (1985); 32, 6238 (1985).
14. I. A. Avrutskii, A. S. Svakhin and V. A. Sychugov, *Opt. Spectrosc. (USSR)* 63, 198 (1987).
15. M. G. Cavalcante, G. A. Farias and A. A. Maradudin, *J. Opt. Soc. Am. B* 4, 1372 (1987).
16. Z. Chen and H. J. Simon, *J. Opt. Soc. Am. B* 5, 1396 (1988).
17. R. Dragila, B. Luther-Davies and S. Vukovic, *Phys. Rev. Lett.* 55, 1117 (1985).
18. T. C. Paulick, *J. Appl. Phys.* 64, 1384 (1988).
19. R. W. Gruhlke, W. R. Holland and D. G. Hall, *Phys. Rev. Lett.* 56, 2838 (1986).
20. R. W. Gruhlke, W. R. Holland and D. G. Hall, *Opt. Lett.* 12, 364 (1987).



21. R. W. Gruhlke and D. G. Hall, Appl. Phys. Lett. 53, 1041 (1988).
22. P. T. Leung, Y. S. Kim and T. F. George, Phys. Rev. B 38, 10032 (1988).
23. Y. S. Kim, P. T. Leung and T. F. George, Chem. Phys. Lett. 152, 453 (1988).
24. A. A. Maradudin and D. L. Mills, Phys. Rev. B 11, 1392 (1975).
25. D. L. Mills and A. A. Maradudin, Phys. Rev. B 12, 2943 (1975).
26. C. T. Tai, Dyadic Green's Functions in Electromagnetic Theory (Intext, San Francisco, 1971).
27. G. S. Agarwal, Phys. Rev. B 14, 846 (1976).
28. E. D. Palik, Ed., Handbook of Optical Constants of Solids (Academic, New York, 1985), p. 313.
29. See, e.g., A. P. Alivisatos, D. H. Waldeck and C. B. Harris, J. Chem. Phys. 82, 541 (1985).
30. P. T. Leung, Y. S. Kim and T. F. George, J Chem. Phys, submitted.

### Figure Captions

1. Configuration of the decay-rate problem at a corrugated thin film.
2. Results for the various reduced decay rates for a film with one grating surface, where we have chosen  $d = 200 \text{ \AA}$ ,  $t = 100 \text{ \AA}$ ,  $\epsilon_1 = 1.0$ ,  $\epsilon_3 = 3.6$ ,  $\zeta_{o_1} = 0$ ,  $\zeta_{o_2} = 50 \text{ \AA}$  and  $Q_2 = 10^{-3} \text{ \AA}^{-1}$ . The solid line represents the total rate, and the dotted and dashed lines represent the contributions from the flat and rough components of the film, respectively.
3. Same as in Fig. 2, but only the total rates for various film thicknesses are shown: solid line,  $t = 100 \text{ \AA}$ ; dotted line  $t = 200 \text{ \AA}$ ; dashed line,  $t \rightarrow \infty$ .
4. Results for a supported ( $\epsilon_1 = 1.0$ ;  $\epsilon_3 = 3.6$ ) grating film, where we have taken  $d = 200 \text{ \AA}$ ,  $t = 100 \text{ \AA}$ ,  $\zeta_{o_1} = \zeta_{o_2} = 5 \text{ \AA}$ ,  $Q_1 = Q_2 = 10^{-3} \text{ \AA}^{-1}$ , and the graphical notations are as given in Fig. 2.
5. Same as in Fig. 4, but only the total rates for various film thicknesses are shown: a,  $t = 100 \text{ \AA}$ ; b,  $t = 150 \text{ \AA}$ ; c,  $t = 200 \text{ \AA}$ ; d,  $t \rightarrow \infty$ .
6. Results for a free-standing film ( $\epsilon_1 = \epsilon_3 = 1.0$ ), where all other parameters and notations are as in Fig. 4.
7. Same as in Fig. 6, but only the total rates for various film thicknesses are shown: solid line,  $t = 100 \text{ \AA}$ ; dotted line,  $t = 200 \text{ \AA}$ ; dashed line,  $t \rightarrow \infty$ .
8. Calculation of the (total) decay-rate spectrum using the parameters given in Ref. 19 with  $t = 500 \text{ \AA}$ . The various graphs correspond to different grating periods ( $\Lambda = 2\pi/Q$ ): a,  $\Lambda = 760 \text{ nm}$ ; b,  $\Lambda = 814 \text{ nm}$ ; c,  $\Lambda = 900 \text{ nm}$ ; d,  $\Lambda = 926 \text{ nm}$ ; e,  $\Lambda = 976 \text{ nm}$ . Note that the logarithmic scale is used for the reduced decay rates.

9. Cross-coupling into the long and short-range surface plasmons for the configuration as in Fig. 8 for thinner film thickness ( $t = 150 \text{ \AA}$ ). Only the cases b (dashed line), d (dotted line) and e (solid line) of Fig. 8 are shown.

Fig. 1




Fig.  
20




Fig 3.




Fig 4




Fig. 5






Fig. 6




Fig. 7




Fig. 8





Fig. 9



TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
		Dr. David L. Nelson Chemistry Division Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. J. E. Jensen  
Hughes Research Laboratory  
3011 Malibu Canyon Road  
Malibu, California 90265

Dr. C. B. Harris  
Department of Chemistry  
University of California  
Berkeley, California 94720

Dr. J. H. Weaver  
Department of Chemical Engineering  
and Materials Science  
University of Minnesota  
Minneapolis, Minnesota 55455

Dr. F. Kutzler  
Department of Chemistry  
Box 5055  
Tennessee Technological University  
Cookeville, Tennessee 38501

Dr. A. Reisman  
Microelectronics Center of North Carolina  
Research Triangle Park, North Carolina  
27709

Dr. D. DiLella  
Chemistry Department  
George Washington University  
Washington D.C. 20052

Dr. M. Grunze  
Laboratory for Surface Science and  
Technology  
University of Maine  
Orono, Maine 04469

Dr. R. Reeves  
Chemistry Department  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. J. Butler  
Naval Research Laboratory  
Code 6115  
Washington D.C. 20375-5000

Dr. Steven M. George  
Stanford University  
Department of Chemistry  
Stanford, CA 94305

Dr. L. Interante  
Chemistry Department  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. Mark Johnson  
Yale University  
Department of Chemistry  
New Haven, CT 06511-8118

Dr. Irvin Heard  
Chemistry and Physics Department  
Lincoln University  
Lincoln University, Pennsylvania 19352

Dr. W. Knauer  
Hughes Research Laboratory  
3011 Malibu Canyon Road  
Malibu, California 90265

Dr. K.J. Klaubunde  
Department of Chemistry  
Kansas State University  
Manhattan, Kansas 66506

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. G. A. Somorjai  
Department of Chemistry  
University of California  
Berkeley, California 94720

Dr. J. Murday  
Naval Research Laboratory  
Code 6170  
Washington, D.C. 20375-5000

Dr. J. B. Hudson  
Materials Division  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. Theodore E. Madey  
Surface Chemistry Section  
Department of Commerce  
National Bureau of Standards  
Washington, D.C. 20234

Dr. J. E. Demuth  
IBM Corporation  
Thomas J. Watson Research Center  
P.O. Box 218  
Yorktown Heights, New York 10598

Dr. M. G. Lagally  
Department of Metallurgical  
and Mining Engineering  
University of Wisconsin  
Madison, Wisconsin 53706

Dr. R. P. Van Duyne  
Chemistry Department  
Northwestern University  
Evanston, Illinois 60637

Dr. J. M. White  
Department of Chemistry  
University of Texas  
Austin, Texas 78712

Dr. D. E. Harrison  
Department of Physics  
Naval Postgraduate School  
Monterey, California 93940

Dr. R. L. Park  
Director, Center of Materials  
Research  
University of Maryland  
College Park, Maryland 20742

Dr. W. T. Peria  
Electrical Engineering Department  
University of Minnesota  
Minneapolis, Minnesota 55455

Dr. Keith H. Johnson  
Department of Metallurgy and  
Materials Science  
Massachusetts Institute of Technology  
Cambridge, Massachusetts 02139

Dr. S. Sibener  
Department of Chemistry  
James Franck Institute  
5640 Ellis Avenue  
Chicago, Illinois 60637

Dr. Arnold Green  
Quantum Surface Dynamics Branch  
Code 3817  
Naval Weapons Center  
China Lake, California 93555

Dr. A. Wold  
Department of Chemistry  
Brown University  
Providence, Rhode Island 02912

Dr. S. L. Bernasek  
Department of Chemistry  
Princeton University  
Princeton, New Jersey 08544

Dr. W. Kohn  
Department of Physics  
University of California, San Diego  
La Jolla, California 92037

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. F. Carter  
Code 6170  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. Richard Colton  
Code 6170  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. Dan Pierce  
National Bureau of Standards  
Optical Physics Division  
Washington, D.C. 20234

Dr. R. Stanley Williams  
Department of Chemistry  
University of California  
Los Angeles, California 90024

Dr. R. P. Messmer  
Materials Characterization Lab.  
General Electric Company  
Schenectady, New York 22217

Dr. Robert Gomer  
Department of Chemistry  
James Franck Institute  
5640 Ellis Avenue  
Chicago, Illinois 60637

Dr. Ronald Lee  
R301  
Naval Surface Weapons Center  
White Oak  
Silver Spring, Maryland 20910

Dr. Paul Schoen  
Code 6190  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. John T. Yates  
Department of Chemistry  
University of Pittsburgh  
Pittsburgh, Pennsylvania 15260

Dr. Richard Greene  
Code 5230  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. L. Kesmodel  
Department of Physics  
Indiana University  
Bloomington, Indiana 47403

Dr. K. C. Janda  
University of Pittsburg  
Chemistry Building  
Pittsburg, PA 15260

Dr. E. A. Irene  
Department of Chemistry  
University of North Carolina  
Chapel Hill, North Carolina 27514

Dr. Adam Heller  
Bell Laboratories  
Murray Hill, New Jersey 07974

Dr. Martin Fleischmann  
Department of Chemistry  
University of Southampton  
Southampton SO9 5NH  
UNITED KINGDOM

Dr. H. Tachikawa  
Chemistry Department  
Jackson State University  
Jackson, Mississippi 39217

Dr. John W. Wilkins  
Cornell University  
Laboratory of Atomic and  
Solid State Physics  
Ithaca, New York 14853



ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. R. G. Wallis  
Department of Physics  
University of California  
Irvine, California 92664

Dr. D. Ramaker  
Chemistry Department  
George Washington University  
Washington, D.C. 20052

Dr. J. C. Hemminger  
Chemistry Department  
University of California  
Irvine, California 92717

Dr. T. F. George  
Chemistry Department  
University of Rochester  
Rochester, New York 14627

Dr. G. Rubloff  
IBM  
Thomas J. Watson Research Center  
P.O. Box 218  
Yorktown Heights, New York 10598

Dr. Horia Metiu  
Chemistry Department  
University of California  
Santa Barbara, California 93106

Dr. W. Goddard  
Department of Chemistry and Chemical  
Engineering  
California Institute of Technology  
Pasadena, California 91125

Dr. P. Hansma  
Department of Physics  
University of California  
Santa Barbara, California 93106

Dr. J. Baldeschwieler  
Department of Chemistry and  
Chemical Engineering  
California Institute of Technology  
Pasadena, California 91125

Dr. J. T. Keiser  
Department of Chemistry  
University of Richmond  
Richmond, Virginia 23173

Dr. R. W. Plummer  
Department of Physics  
University of Pennsylvania  
Philadelphia, Pennsylvania 19104

Dr. E. Yeager  
Department of Chemistry  
Case Western Reserve University  
Cleveland, Ohio 41106

Dr. N. Winograd  
Department of Chemistry  
Pennsylvania State University  
University Park, Pennsylvania 16802

Dr. Roald Hoffmann  
Department of Chemistry  
Cornell University  
Ithaca, New York 14853

Dr. A. Steckl  
Department of Electrical and  
Systems Engineering  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. G.H. Morrison  
Department of Chemistry  
Cornell University  
Ithaca, New York 14853